

First photographic record of albinism in chital (*Axis axis*) from Chitwan National Park, Nepal

Abhinaya Pathak^{1*}, Arjun Tamang², Bishal Sinton³, Sujan Khanal⁴, Bikash Pathak⁵, Prashamsa Paudel⁶

To Cite:

Pathak A, Tamang A, Sinton B, Khanal S, Pathak B, Paudel P. First photographic record of albinism in chital (*Axis axis*) from Chitwan National Park, Nepal. *Discovery* 2023; 59: e16d1018

Author Affiliation:

¹Department of National Parks and Wildlife Conservation, Babarmahal, Kathmandu, Nepal

²Nature Guide Association, Chitwan, Nepal

³Chitwan National Park, Kasara, Chitwan, Nepal

⁴Institute of Forestry, Pokhara Campus, Tribhuvan University

⁵Community Based Anti-Poaching Unit, Chitwan, Nepal

⁶Central Department of Zoology, Kathmandu, Nepal

Contact details

Abhinaya Pathak	abhinayapathak17@gmail.com
Arjun Tamang	tamangarjun123456@gmail.com
Bishal Sinton	bishal.sinton.7@gmail.com
Sujan Khanal	info.sujankhanal@gmail.com
Bikash Pathak	pathakbikash2013@gmail.com
Prashamsa Paudel	prashamsapaudel1@gmail.com

*Corresponding Author

Department of National Parks and Wildlife Conservation, Babarmahal, Kathmandu, Nepal
Email: abhinayapathak17@gmail.com

Peer-Review History

Received: 03 January 2023

Reviewed & Revised: 07/January/2023 to 18/January/2023

Accepted: 21 January 2023

Published: February 2023

Peer-Review Model

External peer-review was done through double-blind method.

Discovery

pISSN 2278-5469; eISSN 2278-5450

URL: <https://www.discoveryjournals.org/discovery>



© The Author(s) 2023. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

ABSTRACT

Chital or spotted deer (*Axis axis*) is one of the important cervid in Chitwan National Park of Nepal. They are easily observed roaming around. However, albino chital is barely found in the park. This study presents the first photographic report of albino chital in buffer zone community forest of the park. Albinism is associated with melanin and rare mutation.

Keywords: chital, Chitwan National Park, buffer zone, albinism, melanin

1. INTRODUCTION

Chital or spotted deer (*Axis axis*) (Kingdom: Animalia, Phylum: Chordata, Order: Mammalia, Class: Cetartiodactyla, Family: Cervidae) are extant in Nepal, India, Bangladesh, Bhutan and Sri Lanka. Chital are found in a variety of habitats, but doesn't prefer habitat that characterizes dense moist (evergreen) forests and open semi-desert or desert (Duckworth et al., 2015). In Nepal, the highest population densities approximately 200 chitals per sq. km were reported from the Bardia National Park (Naess and Andersen, 1993; Moe and Wegge, 1994). In a study from Nepal, a total of 207 line transects covering 372.4 km was assessed for tiger prey-base in Chitwan National Park (henceforth, CNP) which revealed the total number of group observations of chital were 18 in Chure hills whereas 153 in plains with total of 171. Similarly, number of individual chital counted were 262 for Chure hills and 1354 for plains with total count of 1616. Regarded as the most abundant prey ungulate species, tiger prey density comprises 52 chitals per sq. km in CNP (DNPWC, 2020).

2. METHODOLOGY

Study area

CNP is the first protected area of Nepal which spreads over four districts, i.e., Chitwan, Nawalparasi (East of Bardaghat Susta or Nawalpur), Parsa and Makwanpur and three provinces, i.e., Bagmati, Madhesh and Gandaki of Central Nepal. Geographically, it is situated between N 27° 20' 19" to 27° 43' 16" longitude and E 83° 44' 50" to 84° 45' 03" latitude whereas its buffer zone lies between N 27° 28' 23" and 27° 70' 38" longitude and E 83° 83' 98" and 84° 77' 38" latitude. Established on September 20, 1973, CNP was enlisted as a world heritage site under the natural category of UNESCO in 1984. Alongside, it is the first park

in the world to receive Conservation Assured Tiger Standards (CATIS) award in 2015. CNP and its buffer zone possesses both Ramsar site and Important Bird Area (IBA). This park is renowned for last surviving illustration of the natural ecosystems of the Terai region which caters vital habitat for important population of several globally threatened species, such as, Greater One-horned Rhinoceros, Royal Bengal Tiger, Gharial Crocodile and Bengal Florican. Nevertheless, the park boasts 70 species of mammals, more than 576 species of birds, 47 species of reptiles, 55 species of amphibians and 120 species of fishes (CNP, 2015; 2021).




Figure 1 Map of CNP and its Buffer Zone of Nepal with the observation location of adult albino chital.

3. DISCUSSION

Albinism in Animal

Variation in color patterns displayed by mammals is influenced by the presence and distribution of pigmentation in their skin, hair and eyes and is especially related to melanin. A few rare mutations cause chromatic disorders, such as albinism, leucism, piebaldism, hypomelanism and melanism (Lucati & López-Baucells, 2017). Albinism is an inherited condition where the melanin is absent (lack the tyrosinase enzyme), in which individuals do not have any pigmentation on their skin, hair and eyes, skin resulting in snow white body with pink limbs, snout and ears (Menon, 2003; Hayashi & Suzuki, 2018; Van Grouw, 2006; Acevedo et al., 2009). On the contrary, melanism, which is related to dark outer pigmentation is also observed in animals. For instance, Pathak et al., (2021) reported the first photographic record of melanistic leopard in Nepal. Albinism differs among animal groups. It is estimated that true albinos occur with a frequency of about one in 10,000 births (Binkley, 2001).

Apart from hereditary, albinism can be caused by diet, aging, shock, illness, injury and low-quality habitat (Sage, 1962). Albino animals are prone to predation i.e., lack of crypsis around it (Uieda, 2000; Sandoval-Castillo et al., 2006; Acevedo et al., 2009), in addition to a number of pathologies that can arise in association with albinism, such as visual or immunological defects (Abreu et al., 2013). Albino animal has poor eye sight due to lack of eye pigment which affects vision, more difficulty in searching food or avoid danger (Miller, 2005). Further, they are susceptible to sunburn and cancer due to lack of melanin (Halls, 2010). Because of all these reasons, the mating ability is reduced and likely to demise before they transfer the unique genes. Due to their rarity, they are targeted by poachers for illicit trade (Dharmarathne and Wijesinghe, 2020).



Figure 2 Adult albino chital observed at Radha Krishna buffer zone community forest of Chitwan National Park, Nepal during January, 2020. (Photo: © Arjun Tamang)

4. CONCLUSION

To conclude, albinism in chital has been recorded across India (Sayyed et al., 2015; Patel, 2020). Patel, (2020) observed absence of herd or other sympatric ungulates assuming that chital fawn may have rejected from the herd for its unusual appearance. However, in our observation we didn't observe such condition as the albino chital was found with its herd foraging in the forest. Further, some villagers reported that the albino chital was chased and subsequently killed by feral dogs after few days of observation.

Acknowledgements

We express our special thanks to Department of National Parks and Wildlife Conservation and Chitwan National Park for the support to conduct this research. We further extend our gratitude to local communities for providing necessary information about the species and displaying profound interest for its conservation.

Informed consent

Not applicable.

Ethical approval

Chital (*Axis axis*) from Chitwan National Park, Nepal was observed in the study. The Animal ethical guidelines are followed in the study for species observation & identification.

Conflicts of interests

The authors declare that there are no conflicts of interests.

Funding

The study has not received any external funding.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

1. Abreu M, Machado R, Barbieri F, Freitas N, Oliveira L. Anomalous colour in Neo tropical mammals: A review with new records for *Didelphis* sp. (Didelphidae, Didelphimorphia) and *Arctocephalus australis* (Otariidae, Carnivora). *Braz J Biol* 2013; 73:185–194.
2. Acevedo J, Aguayo-lobo A, Torres D. Albino Weddell seal at Cape Shirreff, Livingston Island, Antarctica. *Polar Biol* 2009; 32(8):1239–1243.
3. Binkley SK. Color on, color off. Minnesota Conservation Volunteer 2001.
4. CNP. Management Plan for Chitwan National Park and It's Buffer Zone 2013–2017 2015.
5. CNP. Annual Report of Chitwan National Park and It's Buffer Zone (FY-2077/78) 2021.
6. Dharmarathne WDSC, Wijesinghe N. A review of Albinism records in wild animals across Sri Lanka. *WILDLANKA* 2020; 8(4):206–214.
7. DNPWC. Assessment of Ecological Carrying Capacity of Royal Bengal Tiger in Chitwan-Parsa Complex, Nepal. Department of National Parks and Wildlife Conservation, Babarmahal, Kathmandu 2020.
8. Duckworth JW, Kumar NS, Anwarul Islam M, Sagar Baral H, Timmins R. *Axis axis*. The IUCN Red List of Threatened Species 2015.
9. Halls KM. Albino animals, Minneapolis: Millbrook Press 2010.
10. Hayashi M, Suzuki T. Update on albinism. *Pigmentary Skin Disorders*. Springer Cham 2018; 107–121.
11. Lucati F, López-Baucells A. Chromatic disorders in bats: A review of pigmentation anomalies and the misuse of terms to describe them. *Mamm Rev* 2017; 47:112–123.
12. Menon V. A Field Guide to Indian Mammals. Dorling Kindersley (India) Pvt. Ltd. and Penguin Book India (P) Ltd., Delhi, 2003; 201.
13. Miller J. All about albinism. *Missouri Conservationist* 2005; 66(6). <http://mdc.mo.gov/conmag/2005/06/10.html>. Accessed 3 March 2021.
14. Moe S, Wegge P. Spacing behavior and habitat use of Axis deer (*Axis axis*) in lowland Nepal 1994.
15. Naess KJ, Andersen HJ. Assessing census techniques for wild ungulates in Royal Bardia National Park, Nepal. Thesis, Agricultural University of Norway 1993.
16. Patel DA. Observation of ostracism in an albino Spotted Deer. *Mammal Tales* #16, In: *Zoos Print J* 2020; 35(3):25–26.
17. Pathak A, Thapa S, Khanal S, Paudel P, Magar CT. Record of melanistic Leopard in Nepal. *Mammal Tales* #33, In: *Zoos Print J* 2021; 36(7):12–14.
18. Sage BL. Albinism and melanism in birds. *Br Birds* 1962; 55(6):201–220.
19. Sandoval-Castillo J, Mariano-Melendez E, Villavicencio-Garayzar C. New records of albinism in two elasmobranchs: The tiger shark, *Galeocerdo cuvier* and the giant electric ray, *Narcine entemedor*. *Cybiu* 2006; 30(2):191–192.
20. Sayyed A, Mahabal A, Shrivastava DP. Records of Albinism in Spotted Deer *Axis axis* from India: A Review with addition from Pench National Park, Madhya Pradesh. *Zoos Print J* 2015; 30(10):9.
21. Uieda W. A review of complete albinism in bats with five new cases from Brazil. *Acta Chiropterologica* 2000; 2(1):97–105.
22. Van Grouw H. Not every white bird is an albino: Sense and nonsense about colour aberrations in birds. *Dutch Bird* 2006; 28:79–89.